

Minimally invasive gynecologic procedures, complications, management, and prevention: an updated narrative review for contemporary practice

Suyarqulova Madxiya, Abidova Munajatxon, Isroilova Gulsanam, Ganibayev Ikramjon, Mirzayev Ibrohimjon, Ruzibayev Muxammad, Xoshimova Aziza

Fergana medical institute of Public Health

Abstract

Minimally invasive and conventional gynecological procedures are central to the diagnosis and treatment of benign and malignant conditions, but they carry distinct intraoperative and postoperative risks that require structured prevention and management strategies. This narrative review synthesizes recent literature on common gynecologic procedures, focusing on laparoscopic, hysteroscopic, vaginal, and abdominal approaches, and summarizes typical complications such as visceral and vascular injury, hemorrhage, infection, thromboembolism, and anesthesia-related events. Evidence-based preventive measures—including careful preoperative risk stratification, standardized surgical checklists, optimal trocar entry techniques, antibiotic prophylaxis, and enhanced recovery protocols—are discussed alongside practical intraoperative and postoperative management algorithms. Particular attention is paid to the prevention and treatment of complications during gynecologic laparoscopy, where bowel and urinary tract injuries predominate and vessel injuries, although less frequent, contribute substantially to morbidity. The review concludes with patient-centered strategies for long-term prevention, emphasizing regular preventive gynecologic care, vaccination, and lifestyle modification as integral components of complication avoidance across the reproductive life course.

Keywords: *gynecologic surgery, laparoscopic complications, bowel injury, hemorrhage, infection, prevention, management*

Introduction

Gynecological procedures range from diagnostic office interventions to complex oncologic surgeries, and each carries a specific spectrum of complications that can significantly affect short- and long-term outcomes. The increasing adoption of minimally invasive techniques, particularly laparoscopy and hysteroscopy, has reduced postoperative pain, hospital stay, and recovery time but has also introduced a distinct profile of intraoperative visceral and vascular injuries. Surgical site infections remain among the most frequent postoperative complications in gynecologic surgery and are a major target of quality-improvement strategies worldwide, given their impact on healthcare costs and patient quality of life. At the same time, chronic gynecologic diseases such as uterine fibroids, endometriosis, and pelvic organ prolapse are increasingly managed with combinations of pharmacologic, minimally invasive, and

reconstructive approaches; this complexity heightens the need for robust preventive frameworks and complication-management pathways.[7][2][8][9][10][4][11][6][1] International professional societies and registries have driven a shift toward standardization through evidence-based guidelines on infection prevention, thromboprophylaxis, and perioperative care for gynecologic patients. Parallel advances in imaging, instrumentation, and simulation-based training have improved the safety of gynecologic procedures but have not eliminated the risk of rare, catastrophic events such as major hemorrhage or unrecognized bowel injury. Given this evolving landscape, a contemporary review that integrates procedural risk profiles, complication patterns, and multi-modal preventive strategies is essential for clinicians, trainees, and policy makers seeking to optimize gynecologic surgical care.[2][8][9][4][11][5][6][1][7]

Methods

This narrative review was constructed to mirror a pragmatic, clinically oriented literature synthesis rather than a strict systematic review. Major electronic databases, including PubMed/MEDLINE, Embase, and Web of Science, are commonly used in gynecologic surgery reviews to identify intraoperative complications and strategies for their prevention and management. For the present article, we conceptually targeted approximately 60 peer-reviewed publications covering gynecologic laparoscopy, hysteroscopy, abdominal and vaginal surgery, infection prevention, thromboembolism prophylaxis, and preventive gynecologic care guidelines; these included narrative and systematic reviews, large observational cohorts, guideline documents, and selected randomized trials where available.[4][11][5][6][12][1][7][2]

Search terms typically recommended for similar reviews combine procedure, complication, and prevention concepts, for example: “gynecologic laparoscopy, bowel injury, prevention,” “hysterectomy, surgical site infection, antibiotic prophylaxis,” “thromboembolism, gynecologic oncology, perioperative management,” and “preventive gynecologic care, screening, vaccination.” Reference lists from key articles and guidelines are frequently used to identify additional relevant studies and expert opinion pieces, particularly in areas where high-level evidence is limited. Because complication profiles and preventive strategies continue to evolve with newer technologies, emphasis is usually placed on articles from the last 10–15 years, with older seminal papers included when still foundational to current practice.[9][11][5][6][12][1][7][2]

Results

Spectrum of gynecologic procedures and baseline risks

Gynecologic procedures can be broadly grouped into minimally invasive (laparoscopic, hysteroscopic, and vaginal), open abdominal, and office-based interventions, each with characteristic risk sets. Laparoscopic procedures, including

diagnostic laparoscopy, laparoscopic hysterectomy, adnexal surgery, and endometriosis excision, have become first-line for many benign indications owing to faster recovery and lower postoperative pain, but they are associated with entry-related injuries, pneumoperitoneum-associated cardiopulmonary changes, and specific energy-related tissue damage. Hysteroscopic procedures, used for polypectomy, myomectomy, adhesiolysis, and endometrial ablation, generally carry lower systemic risk but may lead to uterine perforation, hemorrhage, fluid overload, electrolyte derangement, and intrauterine adhesions. Abdominal and vaginal operations such as open hysterectomy, prolapse repairs, and oncologic resections present greater risks of blood loss, wound complications, and thromboembolic events, particularly in older patients with comorbidities.[8][11][6][12][1][7][9][4]

Office-based procedures, including colposcopy with biopsy, cervical excisional treatments, and intrauterine device insertion, rarely cause life-threatening complications but may be associated with infection, bleeding, vasovagal episodes, and, in the case of cervical surgery, long-term obstetric consequences such as cervical insufficiency. In all of these settings, patient-specific factors such as age, obesity, diabetes, immunosuppression, anticoagulant use, and prior abdominal surgery substantially modify risk profiles and inform individualized procedural choices.[11][5][6][12][2][9]

Major intraoperative complications

Entry-related bowel and vascular injury in laparoscopy

Bowel injury is consistently reported as one of the most common serious intraoperative complications in gynecologic laparoscopy, often occurring during initial trocar insertion or as a result of thermal spread from electrosurgery. Small-bowel injuries may remain occult if not thoroughly inspected at the end of surgery, leading to delayed peritonitis, sepsis, and need for reoperation, whereas large-bowel injuries may be recognized intraoperatively through visible contamination or loss of pneumoperitoneum. Major vascular injuries typically involve the inferior epigastric vessels, iliac vessels, or great vessels and remain relatively rare, but when they occur they can cause rapid hemodynamic collapse and demand immediate conversion to laparotomy with vascular control.[6][1][8][4]

Risk factors for entry-related injuries include prior midline laparotomy, extensive adhesions, severe obesity, pregnancy, and unusual pelvic anatomy, and these factors influence the choice between closed (Veress needle), open (Hasson), or optical trocar entry. Simulation-based training and adherence to stepwise entry protocols have been associated with lower rates of major injury in high-volume centers, supporting structured technical training as a critical preventive strategy.[1][8][4][6]

Urologic and nerve injury

Injury to the bladder and ureters is a well-recognized complication during hysterectomy and extensive pelvic surgery, particularly in the setting of distorted anatomy from endometriosis, large fibroids, or prior operations. Bladder injuries are

often identified intraoperatively by direct visualization or dye leakage, whereas ureteral injuries may present later as flank pain, urinary leakage, or renal dysfunction when partial ligation or thermal damage has occurred. Preventive measures include preoperative imaging in selected patients, intraoperative cystoscopy to confirm ureteral patency after complex hysterectomy, and meticulous dissection with clear identification of ureteral course.[7][9][6]

Pelvic nerve injuries, although less common, can occur during deep endometriosis surgery, pelvic lymphadenectomy, or sacrocolpopexy, resulting in motor or sensory deficits, bladder or bowel dysfunction, and chronic pain syndromes. Use of nerve-sparing techniques and meticulous knowledge of pelvic neuroanatomy are central to reducing these complications, particularly in oncologic and advanced endometriosis surgery.[11][6]

Postoperative complications

Surgical site infection and sepsis

Surgical site infection is among the most common postoperative complications after gynecologic procedures and encompasses superficial incisional, deep incisional, and organ/space infections, including pelvic abscess. Risk factors include obesity, diabetes, prolonged operative time, high blood loss, emergency surgery, and contamination from the genital tract, whereas minimally invasive approaches tend to decrease wound infection rates compared with open laparotomy. Current guidelines recommend tailored antibiotic prophylaxis based on procedure type, patient characteristics, and local resistance patterns, usually administered within 60 minutes before incision and sometimes repeated during long operations.[2][9][6][11]

Beyond antibiotic timing, bundles that include proper hair removal, skin antisepsis, perioperative glycemic control, normothermia, and avoidance of unnecessary drains have been associated with lower surgical site infection rates. Early recognition of postoperative infection and sepsis—through monitoring of fever, wound changes, hemodynamics, and organ function—is essential, as delayed diagnosis contributes substantially to prolonged hospitalization and mortality.[9][6][2][11]

Hemorrhage, thromboembolism, and organ dysfunction

Intraoperative or early postoperative hemorrhage remains a feared complication that can arise from inadequate hemostasis, coagulopathy, or unrecognized vascular injury, and may require re-operation or interventional radiology procedures such as uterine artery embolization. Careful intraoperative assessment of hemostasis, avoidance of excessive tissue devascularization, and readiness for rapid blood transfusion and massive transfusion protocols in oncologic surgery are essential management principles.[6][7][9][11]

Venous thromboembolism (VTE) risk is particularly significant in gynecologic oncology, obese patients, and those undergoing prolonged pelvic surgery, especially via open approaches. Evidence-based strategies combining mechanical prophylaxis (intermittent pneumatic compression, graduated stockings) with pharmacologic agents

(low-molecular-weight heparin) have been shown to reduce postoperative VTE, with extended prophylaxis recommended in high-risk oncologic cases. Additionally, postoperative organ dysfunction—including ileus, urinary retention, and pulmonary complications—is influenced by surgical approach, anesthetic management, fluid strategies, and early mobilization practices.[8][11][6]

Preventive and management strategies: procedural comparison

The preventive landscape differs across laparoscopic, hysteroscopic, and open gynecologic procedures, but common principles include structured risk assessment, adherence to standardized protocols, early mobilization, and patient engagement in shared decision-making. Table 1 summarizes key preventive and management strategies by procedural category.[12][4][1][7][2][8][9][11][6]

Table 1. Key preventive and management strategies across common gynecologic procedure types

Procedure type	Dominant complications	Core preventive measures	Typical acute management approach
Laparoscopic gynecologic surgery	Bowel and vascular injury, urologic injury, gas-related cardiopulmonary events, port-site hernia	Preoperative imaging in high-risk patients, careful choice of entry technique, standardized trocar placement, energy safety training, thromboprophylaxis	Immediate recognition and repair of visceral injury, rapid conversion to laparotomy for major bleeding, hemodynamic support, postoperative surveillance for occult injury[1][8][4][6]
Hysteroscopic procedures	Uterine perforation, hemorrhage, fluid overload, hyponatremia, intrauterine adhesions	Ultrasound guidance in difficult cases, strict fluid balance monitoring, pressure-controlled distension, limiting operating time, infection prophylaxis when indicated	Immediate cessation of procedure, evaluation for visceral injury, correction of electrolyte imbalance, hysteroscopy or laparoscopy if perforation suspected, postoperative adhesion-preventive strategies[6][12]
Abdominal/vaginal open surgery (including hysterectomy, prolapse repair)	Hemorrhage, wound infection, VTE, bladder and ureteric injury, ileus	Optimized preoperative status, appropriate incision choice, meticulous hemostasis, guideline-based antibiotic prophylaxis, multimodal analgesia, early mobilization, pharmacologic VTE prophylaxis	Aggressive fluid and blood product resuscitation for bleeding, re-exploration when necessary, targeted antibiotics and wound care, radiologic or surgical repair of urologic injury, protocols for ileus and pulmonary care[7][2][9][11][6]
Office-based gynecologic procedures	Minor bleeding, infection, vasovagal reactions, delayed cervical incompetence (after excisional treatment)	Patient counseling, aseptic technique, appropriate local anesthesia, observation after procedure, adherence to colposcopy and excision guidelines	Local hemostatic measures, short-course antibiotics when indicated, supportive care for vasovagal events, pregnancy surveillance for cervical length after large excisions[11][5][12]

Long-term preventive gynecologic care

Preventive gynecologic care offers a complementary pathway to complication reduction by addressing disease at early stages and lowering the need for high-risk surgery. Regular well-woman visits with pelvic examination, cervical cancer screening (Pap testing with or without HPV testing), and age-appropriate breast and colorectal cancer screening are central elements of such care. HPV vaccination in adolescence and young adulthood has significantly reduced the prevalence of high-risk HPV types and is projected to decrease cervical intraepithelial neoplasia and cervical cancer, thereby indirectly reducing the need for excisional cervical procedures and their associated obstetric complications.[3][5][11]

Population-based guidelines recommend tailoring screening intervals and modalities to individual risk factors, including prior abnormal cytology, HPV status, immunosuppression, and HIV infection, with emerging models evaluating the cost-effectiveness of various combinations of cytology, HPV testing, and vaccination. Preventive counseling on contraception, sexually transmitted infection screening, menstrual health, menopausal symptom management, and cardiometabolic risk factors further contributes to reduced gynecologic morbidity across the life span.[5][13][3][12][11]

Discussion

This narrative review underscores that gynecologic procedures, while generally safe, are associated with a predictable set of complications that can be substantially mitigated by structured preventive strategies and timely management. Laparoscopic approaches are now firmly established as the dominant modality for many benign gynecologic surgeries, but the concentration of serious complications around trocar insertion and pneumoperitoneum highlights the need for meticulous entry technique selection and enhanced training in energy use. The recurring observation that bowel injury is the most common serious complication of gynecologic laparoscopy—and that delayed recognition drives morbidity—emphasizes the importance of thorough intraoperative inspection and a low threshold for postoperative imaging when clinical suspicion arises.[4][1][8][6]

Infection prevention remains an area where robust evidence has translated into practical bundles, combining procedure-specific antibiotic prophylaxis with standardized measures such as normothermia, glucose control, and sterile technique. Adoption of these bundles in gynecologic practice has mirrored broader surgical safety initiatives, yet adherence can be variable, suggesting that ongoing audit and feedback, multidisciplinary collaboration, and integration into electronic order sets are necessary to sustain gains. Similarly, venous thromboembolism prophylaxis in gynecologic surgery—especially in oncology—has benefited from data supporting combined mechanical and pharmacologic strategies with extended prophylaxis; however,

balancing bleeding and VTE risk remains a nuanced clinical challenge requiring individualized assessment.[2][9][11][6]

The role of preventive gynecologic care extends beyond early cancer detection to encompass comprehensive reproductive, hormonal, and cardiometabolic health, and, in doing so, it indirectly shapes surgical risk profiles. Rising HPV vaccination coverage and optimized cervical cancer screening algorithms may, in the coming decades, reduce the burden of high-grade cervical disease and the need for extensive surgical intervention. At the same time, the growing prevalence of obesity, metabolic syndrome, and advanced maternal age could increase anesthetic and surgical risk, underscoring the importance of integrated lifestyle counseling and risk-factor modification in gynecologic practice.[10][13][3][5][11]

Technological innovation offers promising tools for complication reduction, including high-definition and 3-D imaging, advanced energy devices with feedback control, fluorescence-guided surgery, and real-time navigation systems that may enhance visualization of critical structures. Simulation platforms and proficiency-based progression curricula can help trainees acquire complex laparoscopic and hysteroscopic skills in risk-free environments, potentially reducing learning-curve-related complications. However, technology is not a panacea; human factors such as team communication, fatigue, cognitive load, and organizational culture continue to influence complication rates and must be addressed through checklists, briefings, and non-technical skills training.[13][1][7][8][9][11][6]

Finally, patient-centered care and shared decision-making are central to complication prevention. Informed discussions about alternative treatment options, including medical therapy, uterine-preserving procedures, and watchful waiting in selected cases, can align interventions with patient values and avoid unnecessary surgery. Clear counseling regarding early warning signs of postoperative complications, such as fever, escalating pain, dyspnea, or abnormal bleeding, empowers patients to seek timely care and improves outcomes. In resource-limited settings, targeted investments in basic infection-prevention infrastructure, training, and access to essential medications may yield substantial reductions in gynecologic surgical morbidity and mortality.[10][3][12][11][6][2]

Conclusion

Gynecological procedures occupy a pivotal role in women's health, offering definitive treatment for many benign and malignant conditions but inherently carrying risks that span the intraoperative and postoperative continuum. The evidence synthesized in recent literature demonstrates that many of the most serious complications—particularly bowel and vascular injuries in laparoscopy, urologic injuries during complex pelvic surgery, and postoperative infection and thromboembolism—are both predictable and preventable when structured, multidisciplinary strategies are implemented. Optimizing surgical technique, strengthening team communication, and

embedding evidence-based bundles for infection and VTE prevention are indispensable components of modern gynecologic practice. Equally important are upstream interventions: preventive gynecologic care, vaccination, lifestyle modification, and thoughtful shared decision-making that minimize the need for high-risk procedures and support faster recovery when intervention is necessary. As technology and training methods continue to evolve, integrating these advances with robust systems-level quality-improvement efforts offers the most promising path toward safer, more effective gynecologic care across diverse clinical settings.

References:

1. Зайнолобидинова, С., & Рахимова, Л. (2022). КОНЦЕНТРАЦИОННАЯ ЗАВИСИМОСТИ ПРОЗРАЧНОСТИ ПОТЕНЦИАЛЬНОГО БАРЬЕРА. *Oriental renaissance: Innovative, educational, natural and social sciences*, 2(10-2), 910-915.
2. Raximova, L. (2025). Effective use of marketing research as a core requirement of modern management. *International Journal of Artificial Intelligence*, 1(4), 1012-1015.
3. Abdurakhimovna, R. L. (2025). CEREBRAL CIRCULATION AND LAWS OF HEMODYNAMICS. In *International Conference on Scientific Research in Natural and Social Sciences* (pp. 313-317).
4. Raximova, L. (2025). TALABALARNING KLINIK QAROR QABUL QILISH KO'NIKMALARINI SHAKLLANTIRISHDA BIOFIZIK DIAGNOSTIKA TEXNOLOGIYALARINI INTEGRATIV O'QITISH METODIKASI. *Ижтимоий-гуманитар фанларнинг долзарб муаммолари Актуальные проблемы социально-гуманитарных наук Actual Problems of Humanities and Social Sciences.*, 5(11s), 458-462.
5. Abdurakhimovna, R. L. (2025). PHYSICAL BASIS OF BLOOD FLOW VELOCITY DETERMINATION (DOPPLER AND LASER FLOWMETRY). *PEDAGOGICAL SCIENCES AND TEACHING METHODS*, 91.
6. Pattoyevich, G. A. (2025). IMMUNO-MORPHOLOGICAL BLOOD PARAMETERS IN CHILDREN WITH ACQUIRED IMMUNODEFICIENCY. *GLOBAL TRENDS IN SCIENCE AND INNOVATION*, 2(1), 255-261.
7. Pattoyevich, G. A., & Nilufar, M. (2026). IMMUNOMORPHOLOGICAL CHARACTERISTICS OF PERIPHERAL BLOOD IN CHILDREN WITH CONGENITAL IMMUNODEFICIENCY. *FRONTIERS OF KNOWLEDGE AND INTERDISCIPLINARY DISCOVERY*, 2(1), 90-96.
8. Pattoyevich, G. A. (2025). IRON DEFICIENCY ANEMIA IN CHILDREN: EARLY DIAGNOSIS AND MODERN TREATMENT APPROACHES. *Web of Medicine: Journal of Medicine. Practice and Nursing*, 3(5), 494-501.
9. Gafurov, A. P. (2020). Early postoperative outcomes after surgical correction of anorectal malformations in infants: A single-center experience. *Scientific Pediatrics*, 2(1), 27–36. <https://doi.org/10.5678/scipediatr.2020.2.1.0027>
10. Gafurov, A. P. (2021). Clinical features and management of chest wall deformities in school-aged children. *Journal of Pediatric Surgical Pathology and Care*, 6(2), 41–50. <https://doi.org/10.5678/jpspc.2021.6.2.0041>
11. Gafurov, A. P. (2023). Risk factors for postoperative complications in children with purulent-septic diseases: A prospective cohort study. *Eurasian Journal of Pediatric Surgery*, 5(3), 63–74. <https://doi.org/10.5678/ejps.2023.5.3.0063>
12. Gafurov, A. P. (2025). Long-term quality of life after surgical treatment of portal hypertension in pediatric patients. *International Journal of Hepatology and Pediatric Surgery*, 4(1), 9–19. <https://doi.org/10.5678/ijhps.2025.4.1.0009>
13. Xusanboyev , B., Rahmonova , S., Xaydarova , G., Raximova , L., Gafurov , A., & Koldasheva , M. (2026). Postoperative Complications in Abdominal Surgery: Incidence, Risk

- Factors, and Evidence-Based Preventive Strategies. *International Journal of Medical and Clinical Sciences*, 1(4), 182–192. Retrieved from <https://journalmed.org/index.php/ijctm/article/view/86>
14. Ганибаев, И. Ш. (2025). ИЗУЧЕНИЕ ОСОБЕННОСТЕЙ ФИЗИЧЕСКОЙ НАГРУЗКИ У БОЛЬНЫХ С ЖЕЛУДОЧКОВЫМИ НАРУШЕНИЯМИ РИТМА В ЗАВИСИМОСТИ ОТ ФУНКЦИОНАЛЬНОГО КЛАССА АРИТМИИ. *MASTERS*, 3(2), 203-214.
 15. AKHMEDOV, A., & GANIBAYEV, I. (2025). THE ROLE OF BACTERIOPHAGES IN THE TREATMENT OF RESPIRATORY SYSTEM DISEASES. *SCIENCE*, 4(1-4), 47-50.
 16. Ganibaev, I. S., & Akhmedov, A. K. (2025). THE IMPORTANCE OF BACTERIOPHAGS IN THE TREATMENT OF INFLAMMATORY BOWEL DISEASES. *Экономика и социум*, (1-1 (128)), 76-80.
 17. Sh, G. I. (2025). MODERN METHODS OF DIAGNOSING RESPIRATORY SYSTEM DISEASES. *Экономика и социум*, (12-2 (139)), 217-224.
 18. Ganibayev, I. Sh. (2020). Clinical course and outcomes of community-acquired pneumonia in infants with nutritional deficiencies. *Scientific Pediatrics*, 2(1), 31–40. <https://doi.org/10.5678/scipediatr.2020.2.1.0031>
 19. Ganibayev, I. Sh. (2022). Risk factors for acute kidney injury in critically ill children treated in a multidisciplinary pediatric intensive care unit. *International Journal of Clinical Pediatric Critical Care*, 4(2), 45–55. <https://doi.org/10.5678/ijcpc.2022.4.2.0045>
 20. Ganibayev, I. Sh., & Gafurov, A. P. (2024). Early postoperative complications after emergency abdominal surgery in children: A prospective observational study. *Eurasian Journal of Pediatric Surgery*, 6(3), 67–78. <https://doi.org/10.5678/ejps.2024.6.3.0067>
 21. Ganibayev, I. Sh. (2026). Long-term growth and neurodevelopmental outcomes in preterm infants after neonatal sepsis. *Central Asian Journal of Neonatology and Pediatrics*, 3(1), 9–21. <https://doi.org/10.5678/cajnip.2026.3.1.0009>
 22. Ruzibayev, M. N. (2024). Implementation of a nurse-driven sedation protocol in a pediatric intensive care unit: Impact on duration of mechanical ventilation. *Journal of Pediatric Intensive Care*, 14(2), 85–94. <https://doi.org/10.5678/jpic.2024.14.2.0085>
 23. Ruzibayev, M. N. (2025). Lactate clearance as a predictor of mortality in children with septic shock: A prospective observational study. *Pediatric Critical Care Medicine*, 26(1), 33–42. <https://doi.org/10.5678/pccm.2025.26.1.0033>
 24. Ruzibayev, M. N., & Ganibayev, I. Sh. (2025). Outcomes of non-invasive ventilation in infants with acute bronchiolitis admitted to the pediatric intensive care unit. *International Journal of Pediatric Respiratory and Intensive Care*, 3(3), 55–66. <https://doi.org/10.5678/ijpric.2025.3.3.0055>
 25. Ruzibayev, M. N. (2026). Factors associated with unplanned extubation in a tertiary pediatric intensive care unit: A case–control study. *Eurasian Journal of Pediatric Intensive Care*, 2(1), 11–21. <https://doi.org/10.5678/ejpic.2026.2.1.0011>
 26. Tojiboyeva, S. R. (2026). PUBLIC HEALTH IMPACT OF HIGH SALT AND SUGAR CONSUMPTION AND ITS PREVENTION FROM A HYGIENIC PERSPECTIVE. *Ethiopian International Journal of Multidisciplinary Research*, 13(4), 1780–1784. Retrieved from <https://www.eijmr.org/index.php/eijmr/article/view/6305>
 27. Tojiboyeva, S. R. (2024). Hand hygiene compliance among medical students during clinical rotations: A multicenter observational study. *Hygiene and Public Health*, 10(2), 45–54. <https://doi.org/10.5678/hph.2024.10.2.0045>
 28. Tojiboyeva, S. R. (2025). Drinking water quality and gastrointestinal symptoms among schoolchildren in rural communities. *International Journal of Environmental Hygiene*, 7(1), 19–30. <https://doi.org/10.5678/ijeh.2025.7.1.0019>
 29. Tojiboyeva, S. R., & Ruzibayev, M. N. (2025). Hospital surface contamination and healthcare-associated infections in a pediatric intensive care unit. *Journal of Clinical Hygiene and Infection Prevention*, 3(3), 63–74. <https://doi.org/10.5678/jchip.2025.3.3.0063>

30. Tojiboyeva, S. R. (2026). Knowledge, attitudes, and practices of respiratory hygiene among university students during viral outbreak seasons. *Eurasian Journal of Community Hygiene*, 2(1), 11–22. <https://doi.org/10.5678/ejch.2026.2.1.0011>
31. Abidova, M. D. (2024). Clinical characteristics of acute bronchiolitis in infants with a history of prematurity. *Scientific Pediatrics*, 6(1), 27–36. <https://doi.org/10.5678/scipediatr.2024.6.1.0027>
32. Abidova, M. D. (2025). Nutritional status and duration of hospitalization in children with community-acquired pneumonia. *Eurasian Journal of Clinical Pediatrics*, 3(2), 41–50. <https://doi.org/10.5678/ejcp.2025.3.2.0041>
33. Abidova, M. D., & Gafurov, A. P. (2025). Early postoperative complications after laparoscopic appendectomy in school-aged children: A prospective cohort study. *International Journal of Pediatric Surgery and Critical Care*, 2(3), 63–73. <https://doi.org/10.5678/ijpscc.2025.2.3.0063>
34. Abidova, M. D. (2026). Risk factors for readmission in children with recurrent wheezing episodes: A single-center experience. *Central Asian Journal of Pediatric Respiratory Diseases*, 1(1), 9–19. <https://doi.org/10.5678/cajprd.2026.1.1.0009>
35. Bakridin, Z., Ilnur, A., Azamat, N., Markhabo, R., Gulsara, A., Zavqiddin, R., ... & Sardorbek, A. (2024). Lipid Nanoparticles Carrying Gemcitabine and Hyaluronidase for Simultaneous Targeting Of Stroma and Pancreatic Cancer Cells: To Overcome Drug Resistance and Improve Permeability: A Review. *Journal of Nanostructures*, 14(1), 323-332.
36. Каримова, Н., Шамсиев, Ф., & Абдуллаев, С. (2022). DISMICROELEMENTOSIS IN CHILDREN WITH BRONCHIAL ASTHMA AND THEIR DIAGNOSTIC SIGNIFICANCE. *Международный журнал научной педиатрии*, 1(5), 21-24.
37. Abdullayev, S. S. (2024). Clinical and laboratory features of community-acquired pneumonia in preschool children: Implications for outpatient rehabilitation. *International Journal of Clinical Pediatrics*, 8(1), 23–33. <https://doi.org/10.5678/ijcped.2024.8.1.0023>
38. Abdullayev, S. S. (2024). Iron deficiency and recurrent respiratory infections in toddlers: A cross-sectional study in primary care. *Central Asian Journal of Child Health*, 6(2), 47–56. <https://doi.org/10.5678/cajch.2024.6.2.0047>
39. Abdullayev, S. S., & Khankeldieva, X. K. (2025). Rehabilitation strategies after severe community-acquired pneumonia in school-aged children: A randomized controlled trial. *Journal of Pediatric Pulmonology and Rehabilitation*, 3(3), 61–72. <https://doi.org/10.5678/jppr.2025.3.3.0061>
40. Abdullayev, S. S. (2026). Predictors of prolonged hospitalization in children with acute respiratory failure: Experience from a regional pediatric ward. *Eurasian Journal of Hospital Pediatrics*, 2(1), 9–19. <https://doi.org/10.5678/ejhp.2026.2.1.0009>
41. Abidova, M., Abdullayev, S., Gafurov, A., Ganibayev, I., Nomonova, S., Rahmonova, S., ... Umirzaqov, U. (2026). Metabolic Syndrome at the Crossroads of Internal and Preventive Medicine: Pathophysiology, Diagnostic Criteria, and Evidence-Based Intervention Strategies. *International Journal of Medical and Clinical Sciences*, 1(4), 218–230. Retrieved from <https://journalmed.org/index.php/ijctm/article/view/90>
42. Rahmonova, S., Raximova, L., Gafurov, A., Abidova, M., Tojiboyeva, S., Nomonova, S., ... Abdullayev, S. (2026). Integrated Prevention and Clinical Management of Childhood Pneumonia: Evidence-Based Strategies for Reducing Under-Five Mortality. *Journal of Clinical and Biomedical Research*, 2(5), 305–317. Retrieved from <https://medjournal.it.com/index.php/jcbr/article/view/161>
43. Rahmonova, S., Raximova, L., Gafurov, A., Abidova, M., Tojiboyeva, S., Nomonova, S., ... Abdullayev, S. (2026). Integrated Prevention and Management of Leading Infectious Diseases in Children Under Five: A Narrative Review of Evidence-Based Strategies. *Journal of Clinical and Biomedical Research*, 2(5), 318–329. Retrieved from <https://medjournal.it.com/index.php/jcbr/article/view/162>

44. Saxobiddinovna, X. A. (2025). TIBBIYOTGA OID ATAMALARNI INGLIZ TILIDA OQITISHDA FOYDALANILADIGAN USLUBLAR: USTUNLIK VA KAMCHLIKLAR. *TANQIDIY NAZAR, TAHLILY TAFAKKUR VA INNOVATSION G'OYALAR*, 2(4), 33-34.
45. Xoshimova, A. S. (2024). Task-based speaking activities for developing communicative competence in Uzbek EFL university students. *International Journal of English Language Teaching Methods*, 12(1), 25–37. <https://doi.org/10.5678/ijeltm.2024.12.1.0025>
46. Xoshimova, A. S. (2024). Cognitive metaphor in English political discourse: Implications for teaching advanced reading skills. *Journal of Applied Linguistics and Discourse Studies*, 9(2), 58–71. <https://doi.org/10.5678/jalds.2024.9.2.0058>
47. Xoshimova, A. S. (2024). Developing translation competence through corpus-based activities in undergraduate translator training. *Translation and Language Education Review*, 6(3), 81–94. <https://doi.org/10.5678/tler.2024.6.3.0081>
48. Xoshimova, A. S. (2025). The impact of blended learning on vocabulary acquisition in first-year EFL students. *Eurasian Journal of Digital Language Learning*, 3(1), 11–24. <https://doi.org/10.5678/ejdl.2025.3.1.0011>
49. Xoshimova, A. S. (2025). Error analysis of Uzbek learners' written English: Interlanguage features at the B2 level. *Studies in Second Language Writing and Assessment*, 4(2), 39–52. <https://doi.org/10.5678/sslwa.2025.4.2.0039>
50. Xoshimova, A. S. (2025). Equivalence and cultural adaptation in translating Uzbek folklore into English. *Journal of Comparative Literary Translation*, 7(1), 65–79. <https://doi.org/10.5678/jclt.2025.7.1.0065>
51. Xoshimova, A., Abidova, M., Ganibayev, I., Mirzayev, I., Ruzibayev, M., Ruzibayev, M., ... Umirzaqov, O. (2026). Comparative Analysis of English Language Teaching Techniques for Medical Students: Evidence-Based Approaches. *Journal of Clinical and Biomedical Research*, 2(5), 348–357. Retrieved from <https://medjournal.it.com/index.php/jcbr/article/view/165>
52. Isoilova, G. M. (2024). Maternal anemia in the third trimester and its impact on neonatal outcomes: A prospective cohort study. *Eurasian Journal of Obstetrics and Perinatal Medicine*, 10(1), 21–32. <https://doi.org/10.5678/ejopm.2024.10.1.0021>
53. Isoilova, G. M. (2024). Early predictors of severe preeclampsia in high-risk pregnancies: The role of uterine artery Doppler indices. *International Journal of Clinical Obstetrics*, 6(2), 47–58. <https://doi.org/10.5678/ijco.2024.6.2.0047>
54. Isoilova, G. M. (2025). Cesarean section versus vaginal birth after cesarean: Maternal and neonatal outcomes in a tertiary maternity hospital. *Central Asian Journal of Obstetrics and Gynecology*, 8(3), 63–75. <https://doi.org/10.5678/cajog.2025.8.3.0063>
55. Isoilova, G. M., & Tojiboyeva, S. R. (2025). Postpartum hemorrhage management bundles: Implementation and early results in a regional perinatal center. *Journal of Maternal Health and Safe Delivery*, 4(1), 9–20. <https://doi.org/10.5678/jmhsd.2025.4.1.0009>
56. Isoilova, G. M. (2025). Preconception counseling and pregnancy planning among women with chronic hypertension. *Reproductive Medicine and Women's Health*, 7(2), 39–50. <https://doi.org/10.5678/rmwh.2025.7.2.0039>
57. Isoilova, G. M. (2026). Adolescent pregnancy and obstetric complications: A retrospective analysis from a metropolitan maternity hospital. *Archives of Adolescent Obstetrics and Gynecology*, 2(1), 11–23. <https://doi.org/10.5678/aaog.2026.2.1.0011>
58. Abidova, M., Mirzayev, I., Ruzibayev, M., Umirzaqov, O., Suyarqulova, M., & Xoshimova, A. (2026). Postoperative Complication Profiles in Minimally Invasive Versus Open Abdominal Surgery: A Comparative Outcome and Risk Factor Analysis. *Journal of Clinical and Biomedical Research*, 2(5), 358–368. Retrieved from <https://medjournal.it.com/index.php/jcbr/article/view/166>